

Isoëto-Nanojuncetea in Puglia (S-Italy): first phytosociological survey

P. Ernandes¹, D. Gigante², L. Beccarisi³, S. Marchiori⁴, R. Venanzoni², V. Zuccarello⁵

¹Via Salvemini 29, I-73020 Cutrofiano (LE), Italy.

²Department of Chemistry, Biology and Biotechnology, University of Perugia, Perugia, Italy.

³Torre Guaceto State Nature Reserve, Via S. Anna 6, I-72012 Carovigno (BR), Italy.

⁴Via B. Buozzi 21, I-57016 Rosignano M.mo (LI), Italy.

⁵DiSTeBA, University of Salento, Lecce, Italy.

Abstract

Mediterranean temporary ponds (MTP) are considered among the most interesting habitats in the Mediterranean bioclimatic region due to the range of rare and threatened taxa they host. Their ephemeral vegetation presents a particular floristic composition, adapted to unstable ecological conditions due to the alternation between flood and drought. Indeed, these amphibian plant species are established on soils flooded in winter and dry since early summer. The water depth and inundation period determine the plant communities distribution. The vegetation can be referred to the class *Isoëto-Nanojuncetea*, that has a wide distribution and is represented in the study area by two orders: *Isoëtetalia*, Mediterranean and W-European, with spring and early summer bloom, and *Nanocyperetalia*, C-European, with late summer and autumn bloom. This study describes for the first time the *Isoëto-Nanojuncetea* plant communities in Puglia Region, in SE-Italy, where the class distribution was unknown up to now. The here adopted syntaxonomical scheme is consistent with numerical analysis, resulting in homogenous floristic and ecological communities that correspond to syntaxa at association level. The phytosociological approach to the interpretation and management of vegetation and biodiversity has been recently revived by the habitat-type classification adopted by the European 92/43/EEC “Habitat” Directive in its Annex I. The here considered plant communities belong to the priority habitat “3170* Mediterranean temporary ponds”. Aiming to typify them, 170 surveys were carried out in temporary pools over a wide area resulting from a detailed survey that took into account the whole Puglia territory. These complexes of vegetation are very sensitive to the environmental and anthropic disturbance, so they might be good bioindicators with high value for habitat monitoring and conservation.

Key words: amphibious vegetation, biodiversity, “Habitat” Directive 92/43/EEC, Mediterranean, Natura 2000, priority habitat, temporary wetlands.

Introduction

Definition of Mediterranean temporary ponds

Mediterranean temporary ponds (MTP) are freshwater habitats, which are flooded in winter or in the end of spring and dry out in summer with a particular water dynamic that relies mainly on rainfall and diffuse run-off cycle (Grillas *et al.*, 2004). The “Habitats Directive” (92/43/EEC) defines MTP as: “temporary, very shallow water bodies (a few centimetres deep) existing only in winter or at the end of spring, with a semiaquatic Mediterranean vegetation consisting of therophytes and geophytes belonging to the *Isoëtion*, *Nanocyperion flavescens*, *Preslion cervinae*, *Agrostion salmanticae*, *Heleochoilon* and *Lythrion tribracteati* alliances” (European Commission, 2013). With regard to the Italian context, the vegetation typically colonizing these habitat types has been defined as “perennial to annual, amphibious, Mediterranean vegetation, mainly composed by dwarf-sized therophytes and geophytes, with a late winter/early spring phenology, developing in wet systems with shallow temporary ponds, in the Mesosub-, Meso- and Thermo-Mediterranean bioclimatic belts” (Biondi *et al.*, 2012). Also the Ramsar Convention on wetlands, adopted in 1971, devoted the

resolution VIII.33 to temporary ponds defining them as: “small (generally <10 ha), shallow wetlands characterized by alternating phases of drought and flooding and by very self-contained hydrology” (Ramsar Convention Bureau, 2002).

These habitats are found in the five regions of the world with a Mediterranean climate, where their vegetation is characterized by species of the genera *Isoëtes*, *Marsilea* and *Pilularia* (Quezel, 1998; Deil, 2005; Molina, 2005). Temporary ponds are considered to be among the most interesting habitats in the Mediterranean bioclimatic region (Médail *et al.*, 1998; Quézel, 1998) due to the range of rare and critical taxa they host (Biggs *et al.*, 2001; Nicolet *et al.*, 2004).

Ecology of Mediterranean temporary ponds

MTP exhibit great ecological variability depending on their habitat's geology, geomorphology, substrate type, water chemistry and depth, duration of flooding, water source (groundwater or run-off). These environmental features deeply affect also their vegetation, mainly comprising mediterranean therophytic and geophytic species (European Commission, 2013). In general, these communities are characterized by a short life cycle and show a remarkable ability to pro-

duce seeds in the short favourable season, which favours annual plants both in the wet and the dry period. They occur in depressions, which are often endorheic, that are submerged for sufficiently long periods of time to allow the development of hydromorphic soils, aquatic or semi-aquatic vegetation, and specific animal communities (Grillas *et al.*, 2004; Williams, 2006).

Various types of MTP may be distinguished, depending on their origin, the substrate on which they lie, their morphology and their formation. Very different substrate types confer specific physico-chemical characteristics (Gaudillat & Hauray, 2002; Grillas *et al.*, 2004). Deil (2005) distinguishes among waterlogged, amphibic, shallow and deep submerged aquatic conditions. The most characteristic type of MTP develops in topographic depressions on impermeable clayey-marly substrates, or where the water table rises to the surface and floods the ground during the rain season (Grillas *et al.*, 2004).

MTP show a self-regulating hydrology that affects the ecosystem composition (Zacharias & Zamparas, 2010). The length of the hydroperiod and the initiation of the inundation period play a substantial role, depending on the substrates permeability and slope. Changes in hydroperiod have important effects on pattern and strength of biotic interactions such as competition and predation, moreover they represent a criterion for deriving flora and fauna (Spencer *et al.*, 1999; Grillas *et al.*, 2004). The chemical composition of the water is always an important factor for flora and fauna. In addition to the usual physico-chemical characteristics (temperature, pH, dissolved oxygen, electrical conductivity), the ionic composition of the water can affect the presence or abundance of certain species (Chauvelon & Heurieux, 2004).

The spatial and temporal distribution of the vegetation is primarily determined by the water depth gradients and the duration of flooding. The topographical gradients correspond to changes in the duration and depth of flooding and the vegetation is organized principally along these gradients (Rudner, 2005; Grillas *et al.*, 2004).

MTP are therefore good biological indicators in relation to environmental phenomena such as climate changes: for example, the reduction or increase in precipitation, which affect the length of the dry or flooding period, leading to modifications of the hydrological regime and consequently altering the right functioning of these delicate ecosystems (Alvarez Cobelas *et al.*, 2005; Jakob *et al.*, 2003).

The objective of the present study was the investigation of the ephemeral phytocoenoses of Mediterranean temporary ponds located in the Puglia Region, in the territories comprised among the Provinces of Lecce and Brindisi (south Italy).

Materials and Methods

Study area

The present work was carried out in Puglia, that forms the Easternmost part of the Italian peninsula. The territory of Puglia is a long, narrow peninsula, mostly occupied by plains and hills, with some lower mountains belonging to the Southern Apennine and the Gargano Promontory, with high, steep cliffs (Tornadore *et al.*, 1988).

The geological substratum is mainly constituted of limestone in a large platform, with a surface drainage network of modest size, fed only by rainwater (Belmonte, 2000).

From the biogeographic and bioclimatic points of view Puglia located in the Eastern Mediterranean sub-region with a Mediterranean pluviseasonal oceanic bioclimate (Rivas-Martinez *et al.*, 2004a, 2004b), mostly referable to the upper Thermomediterranean and to the upper and lower Mesomediterranean Thermotypes (Pesaresi *et al.*, 2014).

The information related to MTP in Puglia is fragmented and incomplete and most of the locations were unknown up to now: only 7 sites were already indicated for this region (Petrella *et al.*, 2005). Only recently, through new records and specific investigations, the knowledge about these ecosystems has increased, together with the interest to protect the habitat and its rare species (Ernandes *et al.*, 2006, 2007, 2010, 2011; Ernandes & Gigante, 2010; Ernandes, 2011; Ernandes & Marchiori, 2012a, 2012b, 2013).

Data sampling and processing

The field survey was based on Braun-Blanquet's classical approach (Braun-Blanquet, 1964), including the latest methodological updates (Biondi, 2011). We carried out 185 phytosociological relevés in the period 2008-2013 during the winter, spring and summer in the month of January, April and May-June. All the relevés have been performed in 40x40 cm² plots; all of them had no slope, thus slope and exposition are not reported in the tables; localities are listed in Appendix I. In order to capture seasonal changes in species composition and phenology, vegetation sampling was conducted once per season. The description of plant communities followed the phytosociological approach (Braun-Blanquet, 1964; Biondi, 2011). The identification of plant species was based on Pignatti (1982) and Tutin *et al.* (1964-1980). The species nomenclature was updated in accordance with Lucarini *et al.* (2014). Specialist MTP plants, communities and habitats were identified according to the European Manual of Union Habitats (European Commission, 2013), Brullo & Minissale (1998), and Rivas-Martinez *et al.* (2001, 2002).

Data were processed by applying a cluster analysis to produce dendrograms of the carried out relevés,

based on the algorithm of complete linkage (Anderson & Underwood, 1997); hierarchical plant community classification was done using MatEdit software package (MatEdit matrix editor, DBPro, Burba *et al.*, 1992). Plant communities were identified as homogeneous groups in the dendrogram of the surveys. For the syntaxonomic frame we followed the standards proposed by Biondi & Blasi (2013), Biondi *et al.* (2014) and, for some syntaxa, Mucina *et al.* (2016).

The habitats in the study area differ geomorphologically and pedologically and, as a consequence, in aspects concerning the water balance. The vegetation is spatially distributed according to the hydrological regime developing in different communities for different types of ponds. The types here investigated can be classified, according to Ernandes & Marchiori (2013), as: a) cupular pools (Fig. 1) that are small cavities (a few tens of square centimetres or square metres), with very reduced catchment areas created by dissolution on rock layers, with depth less than 2 cm; b) dolines (Fig. 2), *i.e.* depressions created by karstic dissolution and/or subsidence with deep soils, generally with depth between 20-40 cm; c) waterlogged soils (Fig. 3), occurring on impermeable substrates with low hydraulic conductivity (clayey-silty substrates) in slight depressions on the land surface, mostly isolated from the water table and often situated in the clearings of woods, with a soil depth less than 20 cm.

Results

The surveyed vegetation has been referred to the class *Isoëto-Nanojuncetea* Br.-Bl. & Tüxen ex Westhoff, Dijk & Paschier 1946, in which two orders were identified: *Isoëtetalia* Br.-Bl. 1935, best represented in Puglia with 13 communities, and *Nanocyperetalia flavescens* Klika 1935 that includes only one. The detected vegetation types, resulting from the hierar-



Fig. 1 - Cupular pools at Felline (Lecce).



Fig. 2 - Dolina at Laccu Feretru (Sternatia, Lecce).



Fig. 3 - Waterlogged soils at Bosco Preti (Brindisi).

chical classification, are described in the next section, including also a short ecological characterization of the upper rank syntaxa. The dendrogram (fig.4) resulting from cluster analysis show the groups belong to the coenoses described; we have only highlighted the groups related to the new associations and subassociations that will be described in detail in the next paragraph. The A cluster, related to *Junco capitati-Isoëtetum histricis*, is subdivided into the subclusters A1, A2, A3 that correspond respectively to the new subassociations *pleurochaetetosum squarrosae*, *solenopsietosum laurentiae* and *cheiloteletosum chloropii*. The B cluster is related to *Triglochino barrelieri-Isolepidetum cernuae* ass. nova while the C one refers to *Pleurochaeto squarrosae-Isoëtetum todaroanae* ass. nova and its two new subassociations *isoetetosum todaroanii* (C1) and *cheiloteletosum chloropii* (C2). Cluster D brings the vegetation communities of the *Junco pygmaei-Isoëtetum velati* and in particular that of *pilularietosum minutae* subass. nova (D1).

The latest two E and F refer respectively to *Tillaea*




Fig. 4 - Dendrogram from the cluster analysis show the associations named A, B, C, D, E, F and the related subassociations described in the text.

vallantii-*Ptychostometum capillaris* ass. nova and *Eryngium pusillum* dominated community.

Description of the detected communities

Ord.: *ISOËTETALIA* Br.-Bl. 1935

The order brings together several thermophilous and sub-thermophilous micro-associations from the W-Mediterranean, growing on siliceous substrates with acidic to neutral pH, which get already dry in spring (Biondi & Blasi, 2013). Type and duration of the soil flooding are changeable and determine the subdivision into alliances.

All.: *ISOËTION* Br.-Bl. 1935

The alliance groups spring dwarf ephemeral phytocoenoses with a Mediterranean distribution, generally developed on small surfaces flooded for a short period, on limestone soils, rich in *Isoëtes* species: *I. histrix*, *I. gymnocarpa*, *I. iapygia* (now named *I. todaroana*) (Troia & Greuter, 2014; Bagella et al., 2015). These plant communities typically show topographic contacts with the small-size annual dry grasslands belonging to the class *Tuberarietea guttatae* (Br.-Bl. in Br.-Bl., Roussine & Nègre 1952) Rivas Goday & Rivas-Martínez 1963 nom. mut. propos. Rivas-Martínez et al. 2002, with which they often develop a chronological sequence.

In the study area, the waterlogged soils, when affected by trampling, sometimes show the presence of nitrophilous elements (e.g. *Poa annua* or *Plantago coronopus*), while the cupular pools can host taxa from the classes *Saginetea* and *Poetea bulbosae*.

Ass.: *CRASSULO VAILLANTII-PTYCHOSTOMETUM CAPILLARIS* ass. nova (Tab. 1)
(*Holotypus*: Rel. 110, Tab. 1)

Typical community of rocky substrates (cupular pools) that develops on margins of pools with a thickness of few centimeters. Singular is the formation of a moss layer dominated by *Ptychostomum capillare* which covers the limestone base and which, by ensuring a moderate water retention, acts as a substrate for the growth of *Tillaea vallantii* and other microphytes. *P. capillare* is present in all the Italian regions (Aleffi et al., 2008). It has been reported as a companion species in the *Campylopus introflexus*-dominated community recorded by Puglisi et al. (2015) in Lazio region, a bryocoenosis that tends to colonize the outermost zones of the ponds on seasonally moist sandy soil. This is the first time that its role in a plant community was surveyed in Italy.

Ass.: *LYTHRO HYSSOPIFOLIAE-CRASSULETUM VAILLANTII* Bagella, Caria, Filigheddu & Farris 2009 (Tab. 2)

Pioneer amphibious community developing in early spring on cupular pools with a diameter less than 1 m, dominated by *T. vallantii* and other small therophytes of the class *Isoëto-Nanojuncetea* on silty substrate is less than 2 cm depth. The presence of *Polypogon maritimus* in 2 rel-

Tab. 1 - *Crassulo vaillantii-Ptychostometum capillaris* ass. nova (Type: CP = cupular pools; Locality: TP = Torre Pali).

Rel. N.	109	110	113	114	
Cover (%)	100	100	100	100	Presences
Locality	TP	TP	TP	TP	
Type of MTP	CP	CP	CP	CP	
Charact. and diff. <i>taxa</i> of the <i>Crassulo vaillantii-Ptychostometum capillaris</i> ass. nova					
<i>Ptychostomum capillare</i> (Hedw.) D.T. Holyoak	4	5	2a	3	4
<i>Tillaea vaillantii</i> Willd.	1	1	1	3	4
Charact. <i>taxa</i> of the upper units					
<i>Juncus bufonius</i> L.	3	3	2a	2a	4
<i>Juncus capitatus</i> Weigel	2a	2a	2a	1	4
<i>Anagallis arvensis</i> L.	2a	2a	.	.	2
<i>Lythrum hyssopifolia</i> L.	+	2a	.	3	3
Other <i>taxa</i>					
<i>Sedum annuum</i> L.	2a	2a	+	.	3
<i>Sagina apetala</i> Ard.	2a	.	1	2a	3
<i>Draba verna</i> L.	2b	2b	.	.	2
<i>Tripolium pannonicum</i> (Jacq.) Dobrocz. subsp. <i>pannonicum</i>	2a	.	.	.	1
<i>Ochlopoa infirma</i> (Kunth) H. Scholz	1	.	.	.	1
<i>Trifolium arvense</i> L.	.	+	.	.	1

evés could represent a differential role with respect to the Sardinian association described by Bagella *et al.* (2009). A similar community was also described for the rock pools at Crete, in Greece (Bergmeier, 2001) but without any specific syntaxonomical location.

Ass.: *ELATINETUM MACROPODAE* Br. Bl 1931 (Tab. 3)

Monospecific community that starts developing in late spring dominated by *Elatine macropoda*, a stenomediterranean species considered Critically Endangered (CE) in Italy (Conti *et al.*, 1997). This community develops in the center of rock pools where the water depth is generally between 2 and 5 cm and the soil thickness about 2 cm. It develops a spatial contact with the above reported *Lythro-Crassuletum* that colonize the wetter core of the rock pools.

Ass.: *JUNCO CAPITATI-ISOËTETUM HISTRICIS* Br.-Bl. 1935 *TYPICUM* Br.-Bl. 1935 (Tab. 4, Rels. 44-116)

Ephemeral community developing on temporarily flooded soils in open patches of cork-oak woodlands in early spring dominated by therophyte, including several of the class *Tuberarietea*, a typical trait of this vegetation type (Molina, 2005). This community is here referred to the association *Junco capitati-Isoëtetum histricis* Br.-Bl. 1935 described on the basis of one single relevé (Braun-Blanquet, 1935, p. 9) carried out in the surroundings of Meloula, Tunisia. Being the only one reported in the mentioned paper, and being defined by the author himself as "typique" (Braun-Blanquet, 1935, p. 8), this relevé must also be considered the holotype for the typical subassociation. Among the species considered characteristic of this association there are: *Isoëtes histrix*, *Juncus capitatus*, *Cicendia*

Tab. 2 - *Lythro hyssopifoliae-Crassuletum vaillantii* Bagella, Caria, Filigheddu & Farris 2009 (Type: CP = cupular pools; Locality: CB = Contrada Badessa, PC = Palude del Capitano, TP = Torre Pali, LS = La Strea).

Rel. N.	8	55	112	39	14	
Cover (%)	90	10	10	65	85	Presences
Locality	BA	PC	TP	PS	TP	
Type	CP	CP	CP	CP	CP	
Charact. and diff. <i>taxa</i> of the ass. <i>Lythro hyssopifoliae-Crassuletum vaillantii</i> Bagella, Caria, Filigheddu & Farris 2009						
<i>Tillaea vaillantii</i> Willd.	5	2a	2a	4	4	5
<i>Lythrum hyssopifolia</i> L.	.	.	+	+	1	3
<i>Polypogon maritimus</i> Willd.	.	.	.	+	+	2
Charact. <i>taxa</i> of the upper units						
<i>Juncus bufonius</i> L.	+	.	+	.	.	2
<i>Juncus hybridus</i> Brot.	.	.	.	+	2b	2
<i>Gaudinia fragilis</i> (L.) P. Beauv.	+	1
<i>Juncus capitatus</i> Weigel	.	.	+	.	.	1
Other <i>taxa</i>						
<i>Polygonum aviculare</i> L.	+	.	.	.	+	2
<i>Gastidium ventricosum</i> (Gouan) Schinz et Thell.	+	1
<i>Ochlopoa infirma</i> (Kunth) H. Scholz	.	.	+	.	.	1
<i>Sedum caespitosum</i> (Cav.) DC.	+	1
<i>Spergula salina</i> (J. Presl et C. Presl) D. Dietr.	+	1

Tab. 3 - *Elatinetum macropodae* Br. Bl. 1931 (Type: CP = Cupular pools; Locality: TP = Torre Pali, FE = Felline).

Rel. N.	15	18	
Cover (%)	100	90	Presences
Locality	TP	FE	
Type	CP	CP	
Charact. and diff. <i>taxa</i> of the ass. <i>Elatinetum macropodae</i> Br. Bl 1931			
<i>Elatine macropoda</i> Guss.	5	5	2
<i>Lythrum portula</i> (L.) D. A. Webb	2a	.	1

filiformis, *Solenopsis laurentia*. The soils are clayey, 10 to 20 cm deep, flooded with the first rains and dried up in spring.

Together with the characteristic species of the association several moss species grow in this community, playing a significative ecological role in organic enrichment of soils and protection of the sediment from external stresses (radiation, wind, erosion etc.). Very peculiar is also the presence of *I. subinermis*, formerly described as a form of *I. histrix* but recently promoted to the species level, based on karyological and morphological traits observed in Calabrian specimens (Cesca & Peruzzi, 2001).

From data analysis different aspects can be detected, that are spatially contiguous in response of hydrological gradients and light expositions. They are here referred to different subassociations and briefly described hereafter.

Ass.: *JUNCO CAPITATI-ISOËTETUM HISTRICIS* Br.-Bl. 1935 *PLEUROCHETETOSUM SQUARROSAE* subass. nova (Tab. 4, Rels. 91-117) (*Holotypus*: Rel. 96, Tab. 4)

ranean maquis with *Myrtus communis*, *Pistacia lentiscus*, *Cystus salvifolius*, *Cystus monspeliensis*. The community is like a grassy cushion consisting mainly of mosses, in particular *Pleurochaete squarrosa*, on which *I. todaroana* a recently described species (Troia & Raimondo, 2010) grows. This association has an autumnal development and grows on thin deposits of soil in the concavities on calcareous plateaus, where the water supply depends both on run-off and superficial waterflow.

Ass. *PLEUROCHAETO SQUARROSAE-ISOËTETUM TODAROANAE* ass. nova *CHEILOTELETOSUM CHLOROPII* subass. nova (Tab. 5)
(*Holotypus*: Rel. 167, Tab. 5)

The subassociation *cheiloteletosum chloropii* is a community in spatial continuity with the *typicum Pleurochaeto squarrosae-Isoëtetum todaroanae*, developing where the slope increases. It is differentiated by the high constancy of *Cheilotela chloropus*, diagnostic species of the proposed subassociation. In late spring and summer most of the species disappear due to the environmental conditions and the absence of precipitations; very peculiar is the ecological role of the moss species for the development of these communities, as highlighted by Poirion & Barbero (1965) for the description of *Isoëtium* in Costa Azzurra. They act as sponges, retaining water and preventing its rapid flow; they preserve the soil moisture and promote the growth of the other species related to the association. Also Poponessi et al. (2015a, 2015b, 2016a, 2016b), showed the co-occurrence of bryophytes and vascular plants due to the climatic conditions of sites. The community is dominated by plants that appear in late winter, such as *I. todaroana* and *P. squarrosa*, and also geophytes as *R. columnae* and *R. bulbocodium* in addition to annual therophytes.

AGROSTIS POURETII-dominated community (Tab. 6)

It is a community typical of dolines, with shallow soils (<10 cm) but a long inundation period, composed of ephemeral grasses that develop in late spring to summer, on raised areas near the margins of ponds that dry up before the core wet areas.

The relèves were performed in grasslands developed in mosaic with perennial grasses-dominated communities, such as *Carex divisa*, occurring on deeper soils. This community has some floristic affinities with the association *Anthoxanto aristati-Agrostietum salmanticae* described by Biondi & Bagella (2009), however it presents some peculiarities such as the presence of *Carex divisa* subsp. *chaetophylla* with high levels of coverage. For this reason, it is here treated only at the level of community, postponing to a deeper investigation its syntaxonomic attribution.

All. *CICENDIO FILIFORMIS-SOLENOPSIS LAURENTIAE* Brullo & Minissale 1998

This phytosociological alliance, described by Brullo & Minissale (1998), groups the Mediterranean communities which show more remarkable thermo-xerophilous features, compared to the Atlantic alliance *Cicendion* (Rivas Goday in Rivas Goday & Borja, 1961) Br.-Bl. 1967 (Deil, 2005; Bagella et al., 2007, 2009, 2013; Gigante et al., 2007, 2013).

In particular, for the waterlogged soils in Puglia, under this alliance a new association is here described that develops in spring, characterized by the high constancy of *Isolepis cernua*.

Ass. *TRIGLOCHINO BARRELIERI-ISOLEPIDETUM CERNUAE* ass. nova (Tab. 7)
(*Holotypus*: Rel. 13, Tab. 7)

This community is like a small turf with microphytes as *Triglochin barrelieri* and other characteristic taxa of the upper units. This community is localized in temporary pools near the coast with the presence of salty soils. In particularly sub-halophilous places, the relevés result rich in *Polypogon maritimus*. Other frequent species are *Juncus hybridus*, *Lythrum hyssopifolia*, *Lythrum tribracteatum*, sometimes with high cover values. Apart from these taxa, the plant community appears rather species-poor with the only exception of Rel. 173.

All. *PRESLION CERVINAE* Br.-Bl. ex Moor 1937

Annual Mediterranean vegetation developed on largely flooded soils with deep waters, generally wet till the end of spring (Biondi & Blasi, 2013). As concerns its ecology, this alliance presents an intermediate character between *Isoëto-Nanojuncetea* and *Phragmito-Magnocaricetea*. It has a W-Mediterranean character and a variable affinity for the water level: inundated during the winter and very dry in summer. In the study areas, the communities belonging to *Preslion cervinae* seem to be linked to palustrine sites, with high water levels and waterlogged soils until late spring.

Ass. *JUNCO PYGMAEI - ISOËTETUM VELATI TYPICUM* (Tab. 8, Rels. 180-181) and *PILULARIETOSUM MINUTAE* subass. nova (Tab. 8, Rels. 178-183)
(*Holotypus*: Rel. 178, Tab. 8)

Amphibious ephemeral vegetation with a Stenomediterranean-Atlantic distribution, characteristic of small depressions on clayey soils with a long inundation period. This community frequently develops along the banks of ponds with extended periods of flooding and water stagnation. The association has been described by Rivas-Goday (1956) in the surroundings of Guadarrama (Embalse de Santillana). The subassociation *typicum* was described later by the same author (Rivas-Goday, 1970).

As concerns the here reported plant community, the

Tab. 5 - *Pleurochaeto squarrosae-Isoëtetum todaroanae* ass. nova, *isoëtetosum todaroanae* subass. nova and *cheiloteletosum chloropii* subass. nova (Type: CP = Cupular pools; Locality: DC = M. Don Cesare, LS = La Strea).

Rel. N.	101*	102	156	103	105	106	107	108	6	7	171	11	12	170	167*	168	104	157	Presences
Cover (%)	100	100	100	100	100	100	100	##	100	100	100	100	95	154.5	100	100	100	90	
Locality	MD	MD	MD	MD	MD	MD	PS	PS	MD	MD	MD	MD	MD	MV	MV	MD	MD	MD	
Type	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	
Charat. and diff. <i>taxa</i> of <i>Pleurochaeto squarrosae-Isoëtetum todaroanae</i> ass. nova, <i>isoëtetosum todaroanae</i> subass. nova																			
Isoëtes todaroana Troia et Raimondo	2b	2a	.	2a	2a	2a	2b	2b	1	1	2a	3	3	2a	2b	1	2b	+	17
Pleurochaete squarrosa (Brid.) Lindb.	5	5	5	5	5	5	5	5	.	.	5	.	.	2a	2b	4	.	.	12
Diff. <i>taxa</i> of the ass. <i>Pleurochaeto squarrosae-Isoëtetum todaroanae</i> ass. nova, <i>cheiloteletosum chloropii</i> subass. nova																			
Cheilotela chloropus (Brid.) Broth.	2a	5	5	2b	4	4	6
Charact. <i>taxa</i> of the upper units																			
Romulea bulbocodium (L.) Sebast. et Mauri	2a	2a	2b	2a	2a	1	2a	1	+	2a	1	.	.	.	1	+	2a	1	15
Polypogon maritimus Willd.	1	2a	2a	+	1	.	+	+	2b	+	3	1	11
Romulea columnae Sebast. et Mauri	2b	2a	2a	2a	.	.	1	.	1	.	1	+	.	1	9
Ranunculus sardous Crantz	+	.	1	.	1	3	2b	2a	2b	5	8
Juncus bufonius L.	+	1	.	.	1	.	.	2a	2b	+	.	+	7
Gaudinia fragilis (L.) P. Beauv.	1	.	1	1	2a	.	2b	.	.	.	5
Lythrum hyssopifolia L.	.	.	+	2a	.	1	+	4
Lotus angustissimus L.	+	+	1
Ranunculus paludosus Poir.	1	2a	1	.	3
Mentha pulegium L.	3	3	2
Oenanthe pimpinelloides L.	+	1
Juncus hybridus Brot.	2m	1
Anagallis arvensis L. subsp. parviflora (Hoffmanns. et Link) Arcang.	1	.	.	.	1
Anagallis minima (L.) E.H.L. Krause	+	.	.	1
Transgressive <i>taxa</i> from the class <i>Helianthemetea</i>																			
Euphorbia exigua L.	1	1	2b	1	1	.	1	+	.	.	1	1	.	1	3	2b	.	.	12
Antoxantum aristatum Boiss.	2a	2a	1	1	2a	+	6
Catapodium rigidum (L.) C.E. Hubb. ex Dony	2a	1	.	.	+	2a	5
Trifolium cherlerii L.	.	.	.	2b	2b	.	.	1	3
Hypochoeris achyrophorus L.	1	.	.	1	+	.	.	.	3
Plantago bellardii All.	3	.	.	1	2
Vulpia ciliata Dumort.	1	1	2
Trifolium scabrum L. subsp. lucanicum (Gasp. ex Gastridium ventricosum (Gouan) Schinz et Theil.	+	+	2
Asterolinon linum-stellatum (L.) Duby	+	1
Cynosurus echinatus L.	+	1
Ornithopus compressus L.	+	.	.	.	1
Other <i>taxa</i>																			
Ranunculus garganicus Ten.	1	2b	2a	2a	2a	1	2b	2b	.	.	+	+	+	+	.	.	1	.	13
Linum bienne Mill.	1	.	.	.	1	2a	1	1	.	.	1	+	1	+	2a	1	.	1	12
Trifolium resupinatum L.	2a	2b	3	2a	2b	2b	2b	4	+	1	10
Bellis annua L.	2b	.	2a	2a	.	.	.	2a	2b	1	1	.	.	1	.	.	1	.	9
Anagallis arvensis L.	1	.	.	+	1	.	.	2a	.	.	.	1	+	+	.	.	.	1	8
Plantago lagopus L.	.	.	.	3	2a	2a	1	+	.	.	.	+	3	.	.	.	2a	.	8
Sedum rupestre L.	1	1	+	1	.	+	.	.	.	1	6
Ornithogalum gussonei Ten.	1	+	+	1	4
Trifolium stellatum L.	2a	.	1	+	.	.	+	.	4
Scandix pecten-veneris L.	+	+	.	r	1	.	.	4
Crepis rubra L.	1	1	3
Hainardia cylindrica (Willd.) Greuter	1	1	.	+	3
Sherardia arvensis L.	+	.	+	.	.	+	.	3
Cerastium pumilum Curtis	+	.	.	1	2a	.	.	3
Carex flacca Schreb.	2b	+	2
Saxifraga tridactylites L.	1	.	.	.	1	2
Triglochin bulbosum L. subsp. barrelieri (Loisel.) Rouy	1	2a	2
Vulpia muralis (Kunth) Nees	2b	.	.	1	2
Trifolium lappaceum L.	2b	2a	2
Cerastium diffusum Pers. subsp. diffusum	2a	+	2
Sideritis romana L.	+	+	2
Ochlopoa infirma (Kunth) H. Scholz	3	.	2a	2
Parentucellia latifolia (L.) Caruel	r	.	+	2
Centaureum pulchellum (Sw.) Druce	1	+	2
Sagina apetala Ard.	1	r	.	.	.	2
Aira caryophyllea L.	+	+	.	.	2

last 2 relevés in Tab. 8 (180 and 181) can be referred to *Junco pygmaei-Isoëtetum velatae* Rivas Goday 1956 *typicum*. The other relevés (178-183) are locally differentiated by the presence of *Pilularia minuta*, which

is here proposed as a diagnostic taxon for the new sub-association *pilularietosum minutae*. A similar community has been observed by Lorenzoni & Paradis (1997) in Corsica.

Ord.: *NANOCYPERETALIA* Klika 1935

This order groups together the plant communities developed in summer-autumn on flooded substrates which start to get dry not before the end of spring, with eutrophic to hypertrophic, subalkaline to basic soils, often salt- and nitrogen-tolerant, with a C-European and Atlantic distribution, also occurring in Mediterranean areas (Biondi & Blasi, 2013).

Tab. 6 - *Agrostis pourretii* (Willd)-dominated community (Type: D = dolines; Locality: LC = Lago del Capraro, LF = Laccu Feretru).

Rel. N.	70	24	71	72	Presences
Cover (%)	100	100	100	100	
Locality	LC	LF	LC	LC	
Type	D	D	D	D	
Charact. taxa of the <i>Agrostis pourretii</i> -dominated community					
<i>Agrostis pourretii</i> Willd.	5	5	4	4	4
Charact. taxa of the upper units					
<i>Lotus angustissimus</i> L.	4	2b	1	+	4
<i>Ranunculus sardous</i> Crantz	.	+	1	.	2
<i>Centaurium pulchellum</i> (Sw.) Druce	.	+	.	.	1
<i>Centaurium maritimum</i> (L.) Fritsch	.	+	.	.	1
Other taxa					
<i>Polypogon maritimus</i> Willd.	1	+	1	.	3
<i>Carex divisa</i> Huds.	.	4	5	5	3
<i>Aira cupaniana</i> Guss.	1	.	.	1	2
<i>Trifolium nigrescens</i> Viv. subsp. <i>nigrescens</i>	1	.	.	.	1
<i>Phalaris minor</i> Retz.	.	1	.	.	1
<i>Symphotrichum squamatum</i> (Spreng.) G.L. Nesom	.	+	.	.	1
<i>Poa sylvicola</i> Guss.	.	.	1	.	1
<i>Rumex pulcher</i> L.	.	.	+	.	1

Tab. 7 - *Triglochino barrelieri-Isolepidetum cernuae* ass. nova (Type: CP = cupular pools, WS = waterlogged soils; Locality: LS = La Strea, BP = Bosco Preti).

Rel. N.	3	5	173	12	14	13*	10	17	16	Presences
Cover (%)	100	85	95	70	70	75	100	100	90	
Locality	PS	PS	BP	PS	PS	PS	PS	PS	PS	
Type	CP	CP	WS	CP	CP	CP	CP	CP	CP	
Charact. e diff. taxa of the ass. <i>Triglochino barrelieri-Isolepidetum cernuae</i> ass. nova										
<i>Isolepis cernua</i> (Vahl) Roem. et Schult.	5	4	4	2a	1	3	5	3	2b	9
<i>Lythrum hyssopifolia</i> L.	r	.	+	.	1	1	4	4	2b	7
<i>Polypogon maritimus</i> Willd.	.	.	.	+	1	2b	1	+	3	6
<i>Juncus hybridus</i> Brot.	2a	2b	.	.	1	1	.	.	2a	5
<i>Triglochin barrelieri</i> Loisel.	.	.	.	+	.	1	1	+	2a	5
<i>Lythrum tribracteatum</i> Salzm. ex Spreng.	.	.	.	4	4	2b	1	.	.	4
Charatt. taxa of the upper units										
<i>Centaurium maritimum</i> (L.) Fritsch	2b	1	2
<i>Centaurium pulchellum</i> (Sw.) Druce	+	1
<i>Bellis annua</i> L.	.	+	1
<i>Antinoria agrostidea</i> (DC.) Parl.	.	.	2b	1
<i>Juncus bufonius</i> L.	.	.	2a	1
<i>Romulea columnae</i> Sebast. et Mauri	.	.	1	1
<i>Juncus pygmaeus</i> Rich. ex Thuill.	.	.	1	1
<i>Agrostis pourretii</i> Willd.	.	.	1	1
<i>Isoetes histrix</i> Bory	.	.	1	1
<i>Isoetes subinermis</i> (Durieu) Cesca et Peruzzi	.	.	1	1
<i>Ranunculus paludosus</i> Poir.	.	.	+	1
<i>Oenanthe pimpinelloides</i> L.	.	.	+	1
Other taxa										
<i>Spergula salina</i> (J. Presl et C. Presl) D. Dietr.	.	.	.	+	+	.	.	.	+	3
<i>Plantago crassifolia</i> Forssk.	1	+	2
<i>Symphotrichum squamatum</i> (Spreng.) G.L. Nesom	1	1	.	.	2

All. *VERBENION SUPINAE* Slavnić 1951

Sub-halophilous nitrophilous Mediterranean phyto-coenoses with an autumnal phenology, developed on long-flooded substrates, generally dominated by prostrate species of various size (Biondi & Blasi, 2013).

ERYNGIUM PUSILLUM-dominated community (Tab. 9)

This plant community develops on very deep soils (> 20cm), and presents two variants, differentiated by the helophytes *Eleocharis multicaulis* and *E. palustris*. It has a late spring-summer cycle, typical of the MTP named doline. It is dominated by *Eryngium pusillum* (syn.: *E. barrelieri* Boiss.), a stenomediterranean

Tab. 8 - *Juncus pygmaei-Isoëtetum velati* Rivas Goday 1956 *typicum* Rivas Goday 1970 (Rels. 180-181) and *pilularietosum minutae* subass. nova (Rels. 178-183) (Type: D = dolines; Locality: BE = Bellimento).

Rel. N.	178	179	182	183	181	180	Presences
Cover (%)	35	15	70	65	15	35	
Locality	BE	BE	BE	BE	BE	BE	
Type	D	D	D	D	D	D	
Charact. and diff. taxa of the ass. <i>Juncus pygmaei-Isoëtetum velati</i> Rivas Goday 1956							
<i>Isoetes velata</i> A. Braun subsp. <i>velata</i>	2b	1	4	4	1	2a	6
<i>Lythrum thymifolia</i> L.	2a	2a	.	.	2a	2b	4
<i>Juncus pygmaeus</i> Rich. ex Thuill.	+	+	2
Charact. taxa of the subass. <i>pilularietosum minutae</i> subass. nova							
<i>Pilularia minuta</i> Durieu ex A. Braun	1	+	r	1	.	.	4
Charact. taxa of the upper units							
<i>Isolepis cernua</i> (Vahl) Roem. et Schult.	1	1	2	.	1	1	5
<i>Eleocharis multicaulis</i> (Sm.) Sm.	+	.	+	.	.	.	2
<i>Juncus hybridus</i> Brot.	.	+	.	.	.	1	2
Other taxa							
<i>Polypogon subspatheus</i> Req.	.	1	2	.	+	1	4
<i>Cynodon dactylon</i> (L.) Pers.	1	+	2
<i>Symphotrichum squamatum</i> (Spreng.) G.L. Nesom	+	1
<i>Ranunculus peltatus</i> Schrank subsp. <i>baudotii</i> (Godr.) Meikle ex C.D.K. Cook	.	.	1	.	.	.	1
<i>Lepidium coronopus</i> (L.) Al-Shehbaz	+	1

Tab. 9 - *Eryngium pusillum*-dominated community (Type: D = dolines; Locality: LF = Laccu Feretru, BE = Bellimento).

Rel. N.	81	82	20	134	126	127	Presences
Cover (%)	60	85	90	35	15	85	
Type	D	D	D	D	D	D	
Charact. taxa of the <i>Eryngium pusillum</i> -dominated community							
<i>Eryngium pusillum</i> Desf.	3	5	4	2b	2a	4	6
Charact. taxa of the upper units							
<i>Ranunculus sardous</i> Crantz	1	.	+	.	.	.	2
<i>Lythrum junceum</i> Banks et Sol.	.	.	.	1	.	.	1
<i>Isolepis cernua</i> (Vahl) Roem. et Schult.	.	.	.	+	.	.	1
<i>Callitriche brutia</i> Petagna	1	.	1
Other taxa							
<i>Eleocharis palustris</i> (L.) Roem. et Schult.	3	2a	2
<i>Spergula salina</i> (J. Presl et C. Presl) D. Dietr.	.	.	1	+	.	.	2
<i>Polygonum aviculare</i> L.	1	2b	2
<i>Eleocharis multicaulis</i> (Sm.) Sm.	1	2	2
<i>Convolvulus arvensis</i> L.	.	.	3	.	.	.	1
<i>Rumex conglomeratus</i> Murray	.	.	1	.	.	.	1
<i>Polypogon maritimus</i> Willd.	.	.	1	.	.	.	1
<i>Symphotrichum squamatum</i> (Spreng.) G.L. Nesom	.	.	.	2a	.	.	1

hemipterophyte, included in the Puglia Regional Red List as vulnerable (VU). It develops in contact with the deepest area of the pond and reflects the progressive lowering of the water level, when spring arrives; at the same time the community tolerate the emergence for long periods. Singular is the presence of the two above-mentioned variants, occurring in different sites following the allophytic gradient. The variant with *Eleocharis multicaulis* was observed in temporary ponds at Masseria Bellimento, in proximity of the Ionian coast, while the variant with *E. palustris* was found at the pond named Laccu Feretru, in the inland areas of Salento.

Spatial distribution and temporal succession of the plant communities

The essential ecological characteristic of temporary wetlands is the alternation of flooded and dry phases; during each of these phases various environmental factors play an important role in the structure and dynamics of the vegetation. The hydrological regime of the pools depends on a range of variables, such as water level, duration of flooding, soil type and topographical gradient.

We note that in a given pool the spatial and temporal distribution (related to changing seasons) of the vegetation is sometimes determined by the water depth gradients and the duration of flooding (e.g. at Laccu Feretru and Bellimento, that are dolines), while in other cases the driving forces are represented by the topographical gradients and slope (e.g. at Macchie Don Cesare, cupular pool and Bosco Preti, waterlogged soil). At Macchie Don Cesare the slope allows water stagnation only for a short time but the formation of a moss layer keeps the soil wet and allows the growth of the MTP species typical of *Pleurochaeto squarrosae-Isoëtetum todaroanae*.

In the dolines three zones are generally recognized: i) a central zone where communities of aquatic annuals are replaced, in spring, by communities of amphibious annuals, such as *Eryngium barrelieri* and *Mentha pulegium* or, in deeper soil, perennials such as *Eleocharis palustris*, and then in summer by communities of hygrophilous terrestrial annual plants such as *Agrostis pourretii*; ii) an intermediate zone where perennial species form a mosaic with annuals; iii) an outer zone, which dries out more quickly, where both hygrophilous vegetation and more generalist terrestrial species are present.

As concerns the waterlogged soils, where the soil is subject to human disturbance, ponds develop along the borders of Mediterranean maquis and firebreaks; in the more shaded areas there is a moss layer on which the typical species of *Junco capitati-Isoëtetum histricis* grow. In the more exposed sites the subassociation *solenopsietosum laurentiae* develop.

New investigations are under way to define the veg-

etation of new sites, also regarding the bryophyte component. In particular the studies are focusing on an area of Salento characterized by a high concentration of MTP sites and biodiversity of related flora, that includes the territories of Supersano, Cutrofiano, Scorrano and Ruffano in the province of Lecce.

Discussions and conclusions

This is the first paper specifically reporting on MTP vegetation in Puglia. The surveyed communities are referred to four alliances: *Isoëtion*, *Preslion*, *Cicendio-Solenopsis* and *Verbenion*. Seven associations and 2 plant communities were found in the study area, 3 of which are here described for the first time as new association, with 6 new subassociations. The order Isoëtetalia is the most represented in the study area, while the order *Nanocyperetalia*, including the summer vegetation, is only represented by the *Eryngium pusillum*-dominated community.

Additionally, the present paper provides new insights on the role of moss species in the vegetation of Italy, a topic which recently received a vivifying pulse as indicated by a number of specific studies (Puglisi & Privitera, 2012; Poponessi *et al.*, 2015a, 2015b, 2016a, 2016b; Puglisi *et al.*, 2015).

MTP represent a priority habitat (Natura 2000 code: 3170*) listed in Annex I of the Directive 92/43/EC. This vulnerable and unstable habitat exists in areas that, due to their specific characteristics, are under significant human and natural pressures and have become prone to extinction (Bagella *et al.*, 2016).

The conservation value of the studied pools is shown by the richness and rarity of the species they host. The results of our study show that several rare plants at national level are distributed in Puglia, e.g. *E. macropoda* Guss., *E. multicaulis* (Sm.) Sm., *P. minuta* Durieu and *I. velata* A. Braun subsp. *velata* are listed as endangered in Italy (Scoppola & Spampinato, 2005). It should be emphasized that *Pilularia minuta* and *Eleocharis multicaulis* are actually present in only one site in Puglia. At a regional level, 7 species are on the Regional Red List (Conti *et al.*, 1997): *Elatine macropoda*, *Eryngium pusillum*, *Isoëtes histrix*, *Juncus pygmaeus*, *Lythrum thymifolia*, *Lythrum tribracteatum*, *Moenchia mantica*.

Regarding IUCN categories, the recent Red List of the Italian Flora, focusing only on policy species (Rossi *et al.*, 2013, 2014), took into account only a restricted number of the typical MTP species, (e.g. *Pilularia minuta*, considered VU at the national level). We suggest to extend protection to others particularly rare species present in just one or two sites such as *Antinoria agrostidea*, *Elatine alsinastrum*, *Eleocharis multicaulis*, *Juncus hybridus*, *Solenopsis laurentia*, *Verbena supina*, *Isoëtes todaroana*, *Lythrum borysthenicum*.

The species that colonize the MTP are habitat-specif-

ic, rare and difficult to detect. Many of these taxa are poorly understood and little studied because of their small size and short life-cycle, and most of them are classifiable according to the concept of rarity for their abundance, distribution, range area, frequency and phytosociology (Rabinowitz, 1981; Izco, 1998).

Additionally, particular care should be given to the fact that these vegetation types display an intrinsically restrict, scattered and fragmented distribution, demanding for appropriate tools for evaluating area and extent of occurrence, as pointed out by Gigante *et al.* (2016b).

Therefore it is necessary to conduct active and constant monitoring to remove negative effects, preserve biological diversity and restore the sites ecological functions (Bagella *et al.*, 2016; Gigante *et al.*, 2016a). Long-term monitoring of distribution, density and extent of the ponds and their vegetation is indispensable in order to safeguard species and communities of conservation priority.

Syntaxonomic scheme

ISOËTO-NANOJUNCETEA Br.-Bl. & Tüxen ex Westhoff, Dijk & Passchier 1946

ISOËTETALIA Br.-Bl. 1935

Isoëtion Br.-Bl. 1935

Crassulo vaillantii-*Ptychostometum capillaris* ass. nova
Lythro hyssopifoliae-*Crassuletum vaillantii* Bagella, Caria, Filigheddu & Farris 2009

Elatinum macropoda Br. Bl 1931

Junco capitati-*Isoëtetum histricis* Br.-Bl. 1935

typicum Br.-Bl. 1935

pleurochaetetosum squarrosae subass. nova

solenopsietosum laurentiae subass. nova

cheiloteletosum chloropii subass. nova

Pleurochaeto squarrosae-*Isoëtetum todaroanae* ass. nova

isoetetosum todaroanii subass. nova

cheiloteletosum chloropii subass. nova

Preslion cervinae Br.-Bl. ex Moor 1937

Junco pygmaei-*Isoëtetum velatae* Rivas Goday 1956

typicum Rivas Goday 1970

pilularietosum minutae subass. nova

Agrostis pourretii-dominated community

Cicendio filiformis-Solenopsion laurentiae Brullo & Minissale 1998

Triglochino barrelieri-*Isolepidetum cernuae* ass. nova

NANOCYPERETALIA Klika 1935

Verbenion supinae Slavnić 1951

Eryngium pusillum-dominated community

Acknowledgments

The authors wish to thank the anonymous reviewers for the precious suggestions which considerably improved the quality of this contribution. A special

thanks to the prof. Michele Aleffi for the kind collaboration in the determination of the bryophytes and to dott. Luigi Palmisano for the sampling.

References

- Aleffi M., Tacchi R. & Cortini Pedrotti C., 2008. Check-list of the Hornworts, Liverworts and Mosses of Italy. *Bocconea* 22: 1-255.
- Alvarez Cobelas M., Catalan J. & Garcia de Jàlon D., 2005. Impacts on inland aquatic ecosystem. In: Moreno M. (Ed.) *Effects of climate change in Spain*. Madrid, Ministerio de Medio Ambiente.
- Anderson M.J. & Underwood A.J., 1997. Effects of gastropod grazers on recruitment and succession of an estuarine assemblage: a multivariate and univariate approach. *Oecologia* 109: 442-453.
- Bagella S., Aleffi M., Azzella M.M., Bolpagni R., Gigante D., Maneli F., Poponessi S. & Puglisi M., 2016. 3170* Stagni temporanei mediterranei. In: Angelini P., Casella L., Grignetti A., Genovesi P. (Eds.), *Manuali per il monitoraggio di specie e habitat di interesse comunitario (Direttiva 92/43/CEE) in Italia: habitat: 82-83*. ISPRA, Serie Manuali e Linee Guida, 142/2016.
- Bagella S., Caria M.C., Farris E. & Filigheddu R., 2007. Issues related to the classification of Mediterranean temporary wet habitats according with the European Union Habitats Directive. *Fitosociologia* 44 (2) Suppl. 1: 245-249.
- Bagella S., Caria M.C., Farris E. & Filigheddu R., 2009. Phytosociological analysis in Sardinian Mediterranean temporary wet habitats. *Fitosociologia* 46 (1): 11-26.
- Bagella S., Caria M.C., Farris M. & Filigheddu R., 2013. Spatial-time variability and conservation relevance of plant communities in Mediterranean temporary wet habitats: a case study in Sardinia. *Plant Biosyst.* 143 (3): 435-442.
- Bagella S., Peruzzi L., Caria M.C. & Filigheddu R., 2015. Unraveling the taxonomy and nomenclature of the *Isoetes histrix* Bory species complex (*Isoetaceae*, *Lycopodiidae*). *Turkish journal of Botany* 39: 383-387.
- Belmonte G., 2000. *Blu di Puglia - Ambienti e itinerari sommersi intorno al Salento*. Lecce. Conte.
- Bergmeier E., 2001. Seasonal pools in the vegetation of Gavdos (Greece), *in situ* conservation required. *Bocconea* 13: 511-516.
- Biggs J., Fox G., Nicolet P., Whitfield M. & Williams P., 2001. Dangers and opportunities in managing temporary ponds. *Freshwater Forum* 17: 71-80.
- Biondi E., 2011. Phytosociology today: Methodological and conceptual evolution. *Plant Biosyst.* 145 Suppl. 1: 19-29.
- Biondi E. & Bagella S., 2009. Phytosociological analysis in Sardinian Mediterranean temporary wet habitats. *Fitosociologia* 46 (1): 11-26.
- Biondi E. & Blasi C., 2013. *Prodromo della Vegetazione Italiana*. MATTM. Società Botanica Italiana. [available online at www.prodromo-vegetazione-italia.org, accessed on 2017, June10]
- Biondi E., Blasi C., Allegrezza M., Anzellotti I., Azzella M.M., Carli E., Casavecchia S., Copiz R., Del Vico E., Facioni L., Galdenzi D., Gasparri R., Lasen C., Pesaresi S., Poldini L., Sburlino G., Taffetani F., Vagge I., Zitti S. & Zivkovic L. 2014. *Plant communities of Italy: The Vegeta-*

- tion Prodrome. *Plant Biosys.* 148 (4): 728-814.
- Biondi E., Burrascano S., Casavecchia S., Copiz R., Del Vico E., Galdenzi D., Gigante D., Lasen C., Spampinato G., Venanzoni R., Zivkovic L. & Blasi C., 2012. Diagnosis and syntaxonomic interpretation of Annex I Habitats (Dir. 92/43/EEC) in Italy at the alliance level. *Plant Sociology* 49 (1): 5-37.
- Braun Blanquet J., 1935. Un joyau floristique et phytosociologique - "*L'Isoetion*" Méditerranéen. *Comm. Bull. Soc. Etude Sci. Nat. Nimes* 47: 1-23.
- Braun Blanquet J., 1964. *Pflanzensoziologie. Grundzüge der Vegetationskunde*. Ed. 3. Springer Verlag. Wien. 865 pp.
- Bridel E.S., 1819. *Muscologiae Recentiorum Supplementum* 4. Ukert, Gotha. 220 pp.
- Burillo S. & Minissale P., 1998. Considerazioni sintassonomiche sulla Classe *Isoëto-Nanojuncetea*. *Itin. Geobot.* 11: 263-290.
- Burba N., Feoli E., Malaroda M. & Zuccarello V., 1992. Un sistema informativo per la vegetazione. Software per l'archiviazione della vegetazione italiana e per l'elaborazione di tabelle. *Manuale di utilizzo dei programmi CETA Book 2*, Gorizia, 78 pp.
- Cesca G. & Peruzzi L., 2001. *Isoëtes (Lycophytina, Isoetaceae)* with terrestrial habitat in Calabria (Italy). *New karyological and taxonomical data. Fl. Medit.* 11: 303-309.
- Chauvelon P. & Heurteaux P., 2004. Hydro-climatic characteristics. In: Grillas P., Gauthier P., Yavercovsky N., Perennou C. (Eds.), *Mediterranean Temporary pools*. Statione Biologique de la Tour du Valat. Le Sambuc, Arles (France).
- Conti F., Manzi A. & Pedrotti F., 1997. *Liste Rosse Regionali delle Piante d'Italia*. Società Botanica Italiana. WWF Italia.
- Deil U., 2005. A review on habitats, plant traits and vegetation of ephemeral wetlands - a global perspective. *Phytocoenologia* 35 (2-3): 533-705.
- Ernandes P., 2011. Il genere *Isoëtes (Pteridophyta, Lycopsida)*: note tassonomiche, ecologia e distribuzione in Puglia. *Ann. Mus. Civ. Rovereto* 26: 347-358.
- Ernandes P., Beccarisi L., Medagli P. & Zuccarello V., 2006. Note sulle conoscenze floristiche degli "Stagni Temporanei Mediterranei" della Puglia centro-meridionale. *Inf. Bot. Ital.* 38 (1): 185-186.
- Ernandes P., Beccarisi L. & Zuccarello V., 2007. L'habitat prioritario "Stagni Temporanei Mediterranei" in Puglia: nuovi dati distributivi e segnalazioni di specie interessanti. *Inf. Bot. Ital.* 39: 271-279.
- Ernandes P., Beccarisi L. & Zuccarello V., 2010. A new species of *Isoëtes (Isoëtaceae, Pteridophyta)* for the Mediterranean. *Plant Biosys.* 144 (4): 805-813.
- Ernandes P. & Gigante D., 2010. *Isoëtes velata* A. Braun subsp. *velata*, Puglia. In: Marchetti D. (Ed.), *Notule Pteridologiche Italiane VIII*. *Ann. Mus. Civ. Rovereto* 25: 113.
- Ernandes P. & Marchiori S., 2012a. A comparative study of two endemic *Isoëtes* species from South Italy. *ISRN Botany* 2012: 1-7. doi: 10.5402/2012/127250.
- Ernandes P. & Marchiori S., 2012b. The rare water fern *Marsilea strigosa* Willd.: Morphological and anatomical observations concerning a small population in a Mediterranean temporary pond in Puglia. *Plant Biosyst.* 146 (1): 1-6.
- Ernandes P. & Marchiori S., 2013. Mediterranean temporary ponds in Puglia (South Italy): a "joyau floristique" to protect. *Acta Botanica Gallica: Botany Letters* 160 (1): 53-64.
- Ernandes P., Prontera E., Beccarisi L. & Zuccarello V., 2011. Laghi e pozzelle, gli habitat effimeri del Salento: il caso di "Laccu Feretru". *Inf. Bot. Ital.* 43 (1): 17-19.
- European Commission, 2013. *Interpretation Manual of European Union Habitats*. EUR 28. April 2013.
- Gaudillat V. & Haury J., 2002. *Habitats Humides*, vol. 3-7. La documentation française, Paris.
- Gigante D., Attorre F., Venanzoni R., Acosta A.T.R., Agrillo E., Aleffi M. *et al.*, 2016a. A methodological protocol for Annex I Habitats monitoring: the contribution of Vegetation science. *Plant Sociology* 53 (2): 77-87.
- Gigante D., Foggi B., Venanzoni R., Viciani D. & Buffa G., 2016b. Habitats on the grid: The spatial dimension does matter for red-listing. *Journal for Nature Conservation* 32: 1-9.
- Gigante D., Maneli F. & Venanzoni R., 2007. L'ecomosaico degli stagni temporanei nella Piana di Ferretto (Perugia, Italia centrale): un Habitat prioritario della Direttiva 92/43/CEE. *Riv. Idrobiol.* 43 (2004-2007): 148-158. Aracne Ed., Roma.
- Gigante D., Maneli F. & Venanzoni R., 2013. Mediterranean temporary wet systems in inland Central Italy: ecological and phytosociological features. *Plant Sociology* 50 (2): 123-142.
- Grillas P., Gauthier P., Yavercovsky N. & Perennou C., 2004. *Mediterranean Temporary Pools*. Le Sambuc, Arles, France. Statione Biologique de la Tour du Valat.
- Jakob C., Poizat G., Veith M., Seitz A. & Crivelli A.J., 2003. Breeding phenology and larval distribution of Amphibians in a Mediterranean pond network with unpredictable hydrology. *Hydrobiologia* 499: 51-61.
- Izco J., 1998. Types of rarity of plants communities. *J. Veg. Sci.* 9: 641-646.
- Lorenzoni C. & Paradis G., 1997. Description phytosociologique d'une mare temporaire à *Elatine brochonii* dans le sud de la Corse - Département/Région: 20. *Société Botanique du Centre-Ouest, Bull. Soc. bot. Cent.-Ouest N.S.* 28: 21-46.
- Lucarini D., Gigante D., Landucci F., Panfilì E. & Venanzoni R., 2015. The anArchive taxonomic Checklist for Italian botanical data banking and vegetation analysis: theoretical basis and advantages. *Plant Biosyst.* 149 (6): 958-965.
- Médail F., Michaud J., Molina J., Paradis G. & Loisel R., 1998. Conservation de la flore et de la végétation des mares temporaires dulcaquicoles et oligotrophes de France méditerranéenne. *Ecol. Medit.* 24 (2): 119-134.
- Molina J.A., 2005. The vegetation of temporary ponds with Isoetes in the Iberian peninsula. *Phytocoenologia* 35 (2-3): 219-230.
- Mucina L., Bültmann H., Dierßen K., Theurillat J.P., Raus T., Čarni A. *et al.*, 2016. Vegetation of Europe: hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *App. Veg. Sci.* 19 (1): 3-264.
- Nicolet P., Biggs J., Fox G., Hodson M.J., Reynolds C., Whitfield M. & Williams P., 2004. The wetland plant and macroinvertebrate assemblages of temporary ponds in United Kingdom and Wales. *Biol. Conserv.* 120 (2): 261-278.
- Pesaresi S., Galdenzi D., Biondi E. & Casavecchia S., 2014. Bioclimate of Italy: application of the worldwide bioclimatic classification system. *Journal of Maps* 10: 538-553.
- Petrella S., Bulgarini F., Cerfolli F., Polito M. & Teofili C., 2005. *Libro Rosso Degli Habitat d'Italia*. WWF Italia Onlus. Roma.

- Pignatti S., 1982. Flora d'Italia. 3 Voll. Edagricole. Bologna.
- Poirion L. & Barbero M., 1965. Groupements à *Isoetes velata* A. Braun (*Isoetes variabilis* Le Grand). Bull. Soc. Bot. Fr. 112: 436-442.
- Poponessi S., Aleffi M., Bricchi E., Maneli F., Venanzoni R. & Gigante D., 2015a. Role of bryophytes in the vegetation of Mediterranean temporary ponds close to the Temperate bioclimatic border. In: Book of Abstracts: 60. International Symposium on Mediterranean Temporary Ponds "PaulisProject", Sassari 15-17/04/2015. ISBN: 978-88-940864-0-9
- Poponessi S., Aleffi M., Maneli F., Venanzoni R. & Gigante D., 2015b. Seasonal variability of bryophytic vs. vascular species in the vegetation of submediterranean temporary pools in central Italy. In: Abstracts: 151. 110° Congresso SBI, Pavia 14/17-09-2015. ISBN 978-88-85915-16-9
- Poponessi S., Aleffi M., D. Gigante & Venanzoni R., 2016a. Updates on the bryophyte flora of the lowland woods and temporary ponds west of Lake Trasimeno (Central Italy). Fl. Medit. 26: 151-162.
- Poponessi S., Aleffi M., Maneli F., Venanzoni R. & Gigante D., 2016b. Temporary wet ecosystems: the role of bryophytes in a habitat of priority importance for conservation at European level. In: Dragičević, S. (Ed.), Bryophyte Conservation - Towards the new European Red List of Bryophytes. 9th Conference of European Committee for Conservation of Bryophytes. Book of Abstracts: 12-13. Natural History Museum of Montenegro, Podgorica, 26pp.
- Puglisi M., Minissale P., Sciandrello S. & Privitera M., 2015. The bryophyte vegetation of the Mediterranean temporary ponds in Italy. Plant Sociology 52 (2): 69-78.
- Puglisi M. & Privitera M., 2012. A synopsis of the Italian bryophyte vegetation. Cryptogamie, Bryologie 33 (4):357-382.
- Quézel P., 1998. La végétation des mares transitoires à *Isoetes* en région méditerranéenne, intérêt patrimonial et conservation. Ecol. Mediterr. 24 (2): 111-117.
- Rabinowitz D., 1981. Seven forms of rarity. In: Synge H. (Ed.), The biological aspects of rare plant conservation: 205-217. John Wiley & Sons. Chichester.
- Ramsar Convention Bureau, 2002. The Ramsar convention on wetlands. Resolution VIII.33: Guidance for identifying, sustainably managing, and designating temporary pools as Wetlands of International Importance. [available online at: <http://www.ramsar.org/>]
- Rivas Goday S., 1956. Aportaciones a la Fitosociología hispánica (Proyectos de comunidades hispánicas). Nota I. Anales Inst. Bot. Cavanilles 13: 333-422.
- Rivas Goday S., 1970. Revisión de las comunidades hispanas de la clase *Isoeto-Nanojuncetea* Br.-Bl. & Tx. 1943. Anales Inst. Bot. Cavanilles 27: 225-276.
- Rivas-Martínez S., Fernández-González F., Loidi J., Lousa M. & Penas A., 2001. Syntaxonomical Checklist of vascular plant communities of Spain and Portugal to association level. Itin. Geobot. 14: 3-341.
- Rivas-Martínez S., Díaz T.E., Fernández-González F., Izco J., Loidi J., Lousa M. & Penas Á., 2002. Vascular plant communities of Spain and Portugal. Addenda to the syntaxonomical checklist of 2001. Itin. Geobot. 15 (1-2): 5-922.
- Rivas-Martínez S., Penas Á. & Díaz T.E., 2004a. Bioclimatic map of Europe. Cartographic Service, University of Leon, Spain.
- Rivas-Martínez S., Penas Á. & Díaz T.E., 2004b. Biogeographic map of Europe. Cartographic Service, University of Leon, Spain.
- Rossi G., Montagnani C., Gargano D., Peruzzi L., Abeli T., Ravera S., Cogoni A., Fenu G., Magrini S., Gennai M., Foggi B., Wagensommer R.P., Venturella G., Blasi C., Raimondo F.M., Orsenigo S. (Eds.), 2013. Lista Rossa della Flora Italiana. 1. Policy Species e altre specie minacciate. Comitato Italiano IUCN. MATTM.
- Rossi G., Montagnani C., Abeli T., Gargano D., Peruzzi L., Fenu G. et al., 2014. Are Red Lists really useful for plant conservation? The New Red List of the Italian Flora as a case study in the perspective of national conservation policies. Plant Biosyst. 148 (2): 187-190.
- Rudner M., 2005. Environmental patterns and plant communities of the ephemeral wetland vegetation. Phytocoenologia 35: 403-420.
- Scoppola A. & Spampinato G. (Eds.), 2005. Atlante delle specie a rischio d'estinzione. Ministero dell'Ambiente e del Territorio. Direzione per la protezione della natura.
- Spencer M., Blaustein L., Shwartz S. & Cohen J.E., 1999. Species richness and the proportion of predatory animal species in temporary freshwater pools: relationships with habitat size and permanence. Ecology Letters 2: 157-166.
- Tornadore N., Marchiori S. & Marcucci R., 1988. Consistenza floristica e caratteristiche corologiche della flora pugliese. Thalassia Salentina 18: 47-54.
- Troia A. & Greuter W., 2014. A critical conspectus of Italian *Isoetes* (*Isoetaceae*). Plant Biosyst. 148: 13-20.
- Troia A. & Raimondo F.M., 2010. *Isoetes todaroana* (*Isoëtaceae*, *Lycopodiophyta*), a new species from Sicily (Italy). Am. Fern. J. 99 (2009): 238-243.
- Tutin T.G., Heywood V.H., Burges N.A., Moore D., Valentine D.H., Walters S.M. & Webb D.A., 1964-1980. Flora Europea. Voll. 1-5. Cambridge University Press.
- Williams D.D., 2006. The biology of temporary waters. Oxford University Press. Oxford.
- Zacharias I. & Zamparas M., 2010. Mediterranean Temporary Ponds. A disappearing ecosystem. Biodiversity and Conservation 19: 3827-3834.

Appendix I: Sampling site (Provinces are Lecce and Brindisi; Type of ponds D: Doline, CP: Cupular pools, WS: Waterlogged soils; Lithology C: Calcareous, L: Limestone).

Localities	Country	Province	Latitude	Longitude	Altitude (m)	Type of ponds	Lithology
Masseria Bellimento	Nardò	LE	40°11'47"	17°55'22"	4	D	C
Penisola della Strea	Porto Cesareo	LE	40°14'24"	17°54'24"	0.5	CP	C
Macchie Don Cesare	Salve	LE	39°51'36"	18°13'21"	106	CP	C
Torre Pali	Salve	LE	39°50'52"	18°12'47"	43	CP	C
Laccu Feretru	Soletto	LE	40°13'00"	18°11'07"	69	D	L
Bosco del Compare	Brindisi	BR	40°39'27"	17°53'14"	20.5	WS	L
Bosco Preti	Brindisi	BR	40°33'54"	17°52'59"	41.5	WS	L
Lago del Capraro	Sternatia	LE	40°13'99"	18°11'65"	68	D	C
Felline	Alliste	LE	39°55'53"	18°07'15"	38	CP	C
C.da. Badessa	Ostuni	BR	40°43'40"	17°31'20"	275	CP	C
Montevergine	Palmariggi	LE	40°09'52"	18°22'53"	42	CP	C
Palude del Capitano	Nardò	LE	40°12'36"	17°55'32"	12	CP	C